

Dosing

Dosing should be based on child's current weight. Do not guess your child's weight. We will be happy to check your child's weight for you, just call our office. We also encourage all of our parents to have a syringe on hand to measure mL for accurate dosing.

Acetaminophen (Tylenol) Dosing

Acetaminophen infants'/children's oral suspension

Less than 12 pounds - contact our office before use

12-17 pounds – 2.5 mL

18-23 pounds – 3.75mL

24-35 pounds – 5mL

36-47 pounds – 7.5mL

48-59 pounds – 10mL

60-71 pounds - 12.5mL

72-95 pounds – 15mL

Ibuprofen Dosing

Do Not Use For Infants under 6 months of age

Infants' Ibuprofen Drops

Under 12 pounds call the office

12-17 pounds – 1.25mL

18-23 pounds – 1.87mL

Ibuprofen Oral Suspension

24-35 pounds – 5mL

36-47 pounds – 7.5mL

48-59 pounds – 10mL

60-71 pounds - 12.5mL

72-95 pounds – 15mL

Oral Rehydration Solution Recipe

Starch-based solution

1 quart of water

1/2 teaspoon of salt

2 oz. (approx. 1 cup) of baby rice cereal

Sugar-based solution

1 quart of clear water

1/2 teaspoon of salt

8 teaspoons of sugar

Use clean water, if in doubt, use bottled water. Measure ingredients accurately, using measuring spoons. Keep solution in the

refrigerator.